

Early Experience with Laparoscopic Inguinal Hernia Repair

NAEEM MUMTAZ, WAQAR ALAM JAN, SHAHIDULLAH AHMAD, VIQAR ASLAM, SAEEDA KHAN, MUHAMMAD IKRAM

Department of Surgery, Lady Reading Hospital, Peshawar

Correspondence to: Naeem Mumtaz, Assistant Professor, Surgical B Unit, PGMI, Lady Reading Hospital, Peshawar. Email: naeemmumtaz666@hotmail.com, Phone: +923459085341.

ABSTRACT

Objective: To evaluate our experience with Laparoscopic hernia repair with regards to, complications, time to full recovery, return to work, and recurrence.

Methodology: This was a descriptive prospective study, performed in the Surgical B Unit of Lady Reading Hospital, Peshawar from 1st January, 2011 to 1st November 2011. Patients were admitted from the out-patient department. Exclusion criteria included-obstructed/strangulated hernia, patients with previous pelvic surgery, patients not fit for general anaesthesia and children. After written informed consent, either a TEP or TAPP repair was done under general anaesthesia. The course of the procedure was noted for any intraoperative complications, total operating time and conversion to open procedure. Any complications in the post-operative period were also noted. Follow-up was done at 1 week, one month, 4 month and 6 month intervals and any long term complications noted.

Results: A total of 44 patients were included in the study. Male to female ratio was 10:1. Mean age was 37.5 years. Thirty six patients presented with primary hernia while 8 patients had recurrent hernia. In 20 patients, hernia was on the right side while in 16 patients it was on the left side. Eight patients had bilateral hernia. Indirect hernia was seen in 25 patients and direct hernia was present in 14 cases. TEP procedure was done in 8 patients while TAPP procedure was carried out in 36 patients. Major complications encountered were: conversion to open procedure in 4 patients, excessive bleeding in 2 patients, surgical emphysema in 3 patients, significant post-operative pain in 6 patients and port site infection in 2 cases. No recurrence was seen at the end of study.

Conclusion: Laparoscopic inguinal hernia repair can be a routine procedure with results comparable to those of open procedures. It is well suited for recurrences. The visualization of structures is clear and leads to a defect-specific closure.

Keywords: Laparoscopic, Hernia, Inguinal, Mesh, Repair.

INTRODUCTION

The surgical history of inguinal hernias dates back to ancient Egypt. From Bassini's heralding of the modern era to today's mesh-based open and laparoscopic repairs, this history parallels closely the evolution in anatomical understanding and development of the techniques of general surgery.^{1,2}

Accounting for 75% of all abdominal wall hernias, and with a lifetime risk of 27% in men and 3% in women, inguinal hernia repair is one of the most commonly performed surgeries in the world.³ Advances in modern day surgical care has made management of this condition simple and highly effective as hernia surgery is mostly carried out as daycare surgery under local anesthesia. Laparoscopic surgery has made great strides in the last decade to the extent that many of the surgical procedures previously thought of as only

reserved for open surgery, have become routine laparoscopic procedures. Open mesh hernioplasty being simple and highly effective has gained worldwide popularity. But this hasn't stopped surgeons from including this procedure in laparoscopic domain as well. However, Laparoscopic herniorrhaphy hasn't gained widespread popularity and its feasibility as a routine procedure has been questioned. Several studies have demonstrated a definite advantage over open repair with regard to reduced post-operative pain⁴⁻⁶ and earlier return to work and normal activities.^{7,8} The objective of the current study is to evaluate the experience of adopting laparoscopic herniorrhaphy as a routine procedure in a busy tertiary care hospital and assess its outcome in terms of benefits to both the surgeons and the patients.

MATERIAL AND METHODS

This was a descriptive interventional prospective study, conducted in Surgical B unit of Postgraduate Medical institute, Lady Reading hospital, Peshawar. Study duration was from 1st January 2011 to 1st November 2011. Forty four consecutive cases of unilateral or bilateral inguinal hernia were included in the study. Inclusion criteria included age above 18 years, both males and females, primary or recurrent inguinal hernia and unilateral or bilateral inguinal hernia. Exclusion criteria were, patients with obstructed or strangulated hernia, patients in paediatric age group, patients with previous pelvic surgery, patients unfit for general anaesthesia, and those not willing for this procedure. All patients were admitted through outpatient department. A pre-operative workup included detail history, taking, a thorough general and systemic examination, and laboratory investigations like FBC, blood urea and glucose, chest X-ray and ECG for patients above 40 years of age. Patients with comorbid conditions were further evaluated with regards to fitness for general anaesthesia. A written informed consent was taken after explaining the procedure to the patients. General anaesthesia was given to all the patients. All patients received Ceftriaxone in a dose of 1 Gm at the induction of anaesthesia.

Laparoscopic approach was either through TAPP or TEP procedure.

The TAPP procedure was initiated by inserting a 10mm port at the infra-umbilical position followed by inspection of the peritoneal cavity and inguinal anatomy. A 5-mm trocar on the left and a 10mm trocar on right were placed one fingerbreadth below the level of the umbilicus just lateral to the rectus muscle. A transverse peritoneal incision was made from the medial umbilical ligament extending laterally to just above the internal ring. The peritoneum and preperitoneal contents were bluntly dissected from the anterior abdominal wall, exposing the myopectineal orifice. The hernia sac was identified and dissected clear of the contents of spermatic cord. Dissection was carried wide enough to accommodate a 15x15 polypropylene mesh covering the hernia orifice and the post wall of inguinal canal and medially into the space of Ritzius. After the introduction of mesh, it was placed and anchored to Cooper's ligament, superomedially and superolaterally to abdominal wall using tackers. Dissected peritoneal flap was repositioned covering the mesh using a few tackers. The procedure ended by closing the

incisions with deep layers closed with vicryl and skin with fine prolene sutures.

In the TEP approach, an infra-umbilical incision was made slightly lateral to the umbilicus and posterior rectus sheath reached by blunt dissection. A 10 mm trocar was placed and laparoscope introduced. Gas was introduced into the space and dissection was carried out extra-peritoneally upto the pubic symphysis. In some cases a glove finger was used act as a balloon during dissection of the space. A 5-mm trocar was placed 2 finger breadths above the pubis and another 10mm trocar was placed midway between the umbilical port and the lower 5mm trocar. The inguinal anatomy was identified, hernia sac dissected away from cord contents and a preperitoneal space prepared for placement of mesh. A polypropylene mesh of 15x15 size was placed and fixed using tackers.

All patients were monitored for any intra operative anaesthetic and surgical complications. Similarly patients were observed for any complications during their stay in the ward. All patients were followed up at intervals of 1 week, 1 month, 2 month, 4 month and 6 month after their discharge from the unit and any long term complications were recorded on their subsequent follow up visits. All the relevant details of patients including complications were recorded on a specifically designed proforma. Results were analysed at the end of the study using SPSS version 17.

RESULTS

A total of 44 patients were studied. Males were 40 and females 4 with a male to female ratio of 10:1. Minimum age was 20 years and maximum 68 with a mean age of 37.5 years. Different age groups affected are shown in Figure No. 1.




Figure No.1

Inguinal hernia as a primary condition was seen in 36(81.8%) cases while 8(18.2%) patients presented with recurrent inguinal hernia. Right sided inguinal hernia was seen in 20(45.45%) patients while it was on the left side in 16(36.36%) cases. it was found bilateral in 8(18.2%) patients.

Types of hernia encountered are given in figure No.2.




Figure No.2 Types of hernia

Table No.1 Complications

Complication	TEP	TAPP	Total	%age
Port site bleeding	1	1	2	4.54%
Surgical Emphysema	1	2	3	6.81%
Neuralgia/pain	4	2	6	13.63%
Conversion to open procedure	1	3	4	9.09%
Port site infection	2	0	2	4.54%
Scrotal edema	2	0	2	4.54%
Urinary Retention	0	1	1	2.27%
Recurrence	0	0	0	0.0%

Hernia was of indirect type in 29(65.90%) patients with undescended testes in 2 of these cases. It was of direct type in 12(27.27%) patients while in 3(6.81%) cases it was of pantaloon type. All patients were operated electively by a single surgeon and given general anaesthesia. A Transperitonealpreperitoneal (TAPP) procedure was done in 36(81.8%) cases while Totally extra

peritoneal(TEP) repair was done in 8(18.2%) patients. The mean operating time for unilateral inguinal hernia repair was 40 minutes (range 30 to 50) and for bilateral repair, it was 63 minutes (range 50 to 70). Complications observed during the procedure and in the post-operative period are given in table 1.

A mean duration of follow-up of 16 weeks was obtained for 40 patients (>90%). Follow-up data was not available for 4 patients. Work or normal physical activity was resumed at a median of 10 days (range 8 to 12 days) after unilateral repair. After discharge six patients sought treatment elsewhere for minor complaints.

DISCUSSION

Inguinal herniorrhaphy is one of the commonest surgical procedure performed worldwide. In the United States, inguinal herniorrhaphy accounts for approximately 800,000 cases annually.⁹

Laparoscopic inguinal herniorrhaphy was first described in the early 90s by GerandShultz.^{10,11} It gained popularity after the success of laparoscopic cholecystectomy. Much work has been done to establish the role of laparoscopic approach in the management of inguinal hernia and to determine whether open or laparoscopic approach is better. The most recent data available on the percentage of laparoscopic herniorrhaphies among all hernia repairs ranges from 30% in Germany, 14% in USA to only 4% in England and Denmark.¹² As trend in adopting laparoscopic techniques for different procedures grows steadily, controversies still exist regarding the feasibility of laparoscopic herniorrhaphy as the procedure of choice. The 2004 VA study recruited almost 1700 patients and demonstrated lower recurrence rates for the traditional approach.¹³

The Italian multicenter randomised controlled trial on 108 patients in 1998 showed more expensive laparoscopy which also requires general anaesthesia is not justified on a routine basis.¹⁴

The current study, was to evaluate our experience and determine if it can be adopted well in our set of circumstances, where cost of treatment, prevention of complications and early return to work is as important as anywhere else.

This study included patients with different types of hernia. Most authorities believe laparoscopy is indicated only in cases of bilateral repair, recurrent hernia, or selected patients, i.e. patients who undergo other laparoscopic procedures, patients who require

the fastest possible recovery (athletes), and patients not amenable to loco-regional anaesthesia.^{15,16} In particular recurrent hernias are effectively approached from behind, working on unaffected issues, allowing less traumatic and non-lengthy dissection.

The TEP may have some advantages over the TAPP in terms of postoperative pain and reduced potential for intraperitoneal complications but does require a high level of technical skill associated with a considerable learning curve.^{17,18}

All patients received prophylactic antibiotic at induction of anaesthesia. Current data remain controversial, but the most recent meta-analysis supports the use of antibiotic prophylaxis when performing a mesh-based repair.¹⁹

A TAPP procedure was the predominant approach as compared to TEP approach in this study. The reason being the surgeon realizing after initial experience, TAPP being easier to perform without the need for additional expensive instruments and had a shorter learning curve.

Laparoscopic herniorrhaphy shares some complications with the open technique but also has its own set of complications. Some of these problems were encountered early on and were corrected as surgeons became more experienced with the technique. Thus, the incidence of complications has decreased with time.²⁰ The overall complication rate in this study was 20.45% taking in to account minor complications as well like scrotal oedema and minor wound infection which were adequately managed.

Felix et al found a 2.7% complication rate over a 6-year period. 90% of the complications occurred in the first 50% of patients.²¹

The main intraoperative complications encountered in this study was port site bleeding in 2 patients. However this did not lead to abandonment of the procedure as bleeding was adequately controlled. Tetik et al and others found a high incidence of hematomas and suggested that it was due to the more extensive dissection performed with the TAPP and TEP repairs.²² In four patients, the procedure was converted to open repair. In one patient undergoing TEP repair, there was massive development of emphysema while in 2 patients there was inadvertent breach in the peritoneum leading to conversion to open procedure. In one patient pelvic adhesions prevented the surgeon from doing a safe dissection. Complications related to the laparoscopic technique itself, or to the hernia

repair, include vascular injuries or injuries to bowel. A number of early studies showed a higher rate of these injuries. However, no such complication was seen in this study.

The majority of complications are postoperative. Urinary retention, a complication of open repair as well, is reported to be the most common of the patient related problems, with an incidence of 1.3% to 5.8%^{20,22,23}. Only one patient in current study had urinary retention which was adequately managed without long term need for any intervention.

Orchitis has been an infrequent complication found in a small number of by different researchers.^{20,22,23} None of patient in this study showed this complication.

Six patients complained of intractable pain in the immediate post-operative period. It was attributed to transient neuralgia and got settled with analgesia without going in to late phase. Fitzgibbons et al looked at postoperative neuralgia and found that leg pain decreased significantly (from 7% to 1.8%) after surgeons performed 30 cases. Postoperative groin pain remained steady at about 8% and testicular pain at 2%.²⁰

Two patients developed port site infection which got settled with conservative treatment. None of patient had deep seated mesh infection. There are reports of mesh getting rejected years after the laparoscopic procedure.^{24,25}

No recurrence of hernia was observed till the end of study period in this study.

The laparoscopic repair can be done with low recurrence rates (0%–1.1%).^{26,27} Other authors, however, have reported higher recurrence rates of 5% to 20%.^{28,29}

CONCLUSIONS

Laparoscopic inguinal hernia repair can be a routine procedure with results comparable to those of open procedures. It is well suited for recurrences and patients presenting with bilateral herniae. The visualization of structures is clear and leads to a defect-specific closure. The advantages of the laparoscopic approach include

1. The type of hernia is obvious,
2. Trocar placement is identical for any side or hernia type,
3. There is clear visualization of the anatomy.
4. Routine video documentation renders the diagnostic accuracy objective and absolute.

However a considerable amount of skill and infrastructure is required in order to make it a routine procedure.

REFERENCES

1. Lau WY. History of treatment of groin hernia. *World J Surg.* Jun 2002;26(6):748-59.
2. Gray SH, Hawn MT, Itani KM. Surgical progress in inguinal and ventral incisional hernia repair. *SurgClin North Am.* Feb 2008;88(1):17-26.
3. Jenkins JT, O'Dwyer PJ. Inguinal hernias. *BMJ.* Feb 2 2008;336(7638):269-72.
4. Wellwood J, Sculpher MJ, Stoker D, et al. Randomised controlled trial of laparoscopic versus open hernia repair for inguinal hernia: outcome and cost. *Br Med J* 1998; 317:103-10
5. Wright DM, Kennedy A, Baxter JN. Early outcome after open versus extraperitoneal endoscopic tension-free hernioplasty. *Surgery* 1996; 119: 552-7
6. Millikan KW, Kosik ML, Doolas A. A prospective comparison of transabdominalpreperitoneal laparoscopic hernia repair versus traditional open hernia repair in a university setting. *SurgLaparoscEndosc* 1994; 4: 247-53
7. Kiruparan P, Pettit SH. Prospective audit of 200 patients undergoing laparoscopic inguinal hernia repair with followup from 1 to 4 years. *J R CollSurgEdin*1998; 43:13-6
8. Brooks DC. A prospective comparison of laparoscopic and tension-free open herniorrhaphy. *Arch Surg*1994; 129: 361-6
9. Rutkow IM. Demographic and socioeconomic aspects of hernia repair in the United States in 2003. *SurgClin North Am.* Oct 2003;83(5):1045-51.
10. Ger R. The laparoscopic management of groin hernias. *ContempSurg* 1991;39(4):15-9.
11. Schultz L, Graber J, Pietrafitta J, et al. Laser laparoscopic herniorrhaphy: a clinical trial, preliminary results. *J LaparoendoscSurg* 1990;1:41-5.
12. Tatulli F, Chetta G, Caputi A, Mastrototaro P, Ruggieri T. Laparoscopic inguinal hernia repair, Audit of our experience with laparoscopic trans-abdominal pro- peritoneal repair (T.A.P.P.)*ChirurgiaItaliana* 2009;61(1):47-53
13. Neumayer L, Giobbie-Hurder A, Jonasson O, Fittzgibbons R Jr, Dunlop D, Gibbs J, Reda D, Henderson W. Veterans Affairs Cooperative Studies Program 456 Investigators. Open mesh versus laparoscopic mesh repair of inguinal hernia. *N Engl J Med* 2004;18:1819-28.
14. Paganini A M et al. A randomized controlled clinical study of laparoscopic vs. open tension-free inguinal hernia repair. *SurgEndosc* 1998; 12:979-86.
15. Puri V, Fittzgibbons RJ Jr. Laparoscopic vs. conventional tension free inguinal herniorrhaphy: 2005 Society of American Gastrointestinal Endoscopic Surgeons (SAGES) annual meeting debate. *SurgEndosc* 2006; 20:1809-16.
16. Lim M, O'Boyle C J, Royston C M S, Sedman P C. Day case laparoscopic herniorrhaphy. *SurgEndosc* 2006; 20:1453-9.
17. Khoury N. A comparative study of laparoscopic extraperitoneal and transabdominalpreperitonealherniorrhaphy. *J LaparoendoscSurg* 1995; 5: 349-55
18. Liem MS, van Steensel CJ, Boelhouwer RU, Clevers GJ, Meijer WS Vente LS, van Vroonhoven TJ. The learning curve for totally extraperitoneal laparoscopic inguinal hernia repair. *Am J Surg* 1996; 171: 281-5'
19. Sanabria A, Dominguez LC, Valdivieso E, Gomez G. Prophylactic antibiotics for mesh inguinal hernioplasty: a meta-analysis. *Ann Surg.* Mar 2007;245(3):392-6.
20. Fittzgibbons RJ Jr, Camps J, Cornet DA, et al. Laparoscopic inguinal herniorrhaphy. Results of a multicenter trial. *Ann Surg* 1995;221(1):3-13.
21. Felix EL, Harbertson N, Vartanian S. Laparoscopic hernioplasty: significant complications. *SurgEndosc* 1999;13(4):328-31.
22. Tetik C, Arregui ME, Dulucq JL, et al. Complications and recurrences associated with laparoscopic repair of groin hernias. *SurgEndosc* 1994;8:1316-23.
23. Phillips EH, Arregui M, Carroll BJ, et al. Incidence of complications following laparoscopic hernioplasty. *SurgEndosc* 1995;9:16-21.
24. D. Foschi, F. Corsi, P. Cellerino, A. Trabucchi, E. Trabucchi. Late rejection of the mesh after laparoscopic hernia repair. *Surgical Endoscopy* 1998;12(5):455-457.
25. C. Hofbauer, P. V. Andersen, P. Juul and N. Qvist. Late mesh rejection as a complication to transabdominalpreperitoneal laparoscopic

- hernia repair. *Surgical Endoscopy* 1998;12(9):1164-1165
26. Felix EL, Michas CA, McKnight RL. Laparoscopic repair of recurrent hernias. *SurgEndosc* 1995;9(2):135-9.
27. Hawashi A, Thao U, Chapital A. Laparoscopic transabdominalpreperitoneal inguinalhernia repair for recurrent inguinal hernia. *Am Surg* 2002;68(3):303-7.
28. Memon MA, Feliu X, Sallent EF, et al. Laparoscopic repair of recurrent hernias. *SurgEndosc* 1999;13(8):807-10.
29. Beets GL, Dirksen CD, Go PMNYH, et al. Open or laparoscopic preperitoneal meshrepair for recurrent inguinal hernia? *SurgEndosc* 1999;13(4):323-7.