

Bacterial urinary tract infection and antibiotic sensitivity pattern among type II diabetic patients – A cross-sectional study from tertiary care teaching hospital in Central Lahore

Kaneez Fatima¹, Khadija Muneer², Naheed Hashmat³, Sana Humayun⁴, Qaiser Pervaiz⁵, Saim Maqsood⁶

¹Woman Medical Officer, GMH Chohan Road, Lahore, ²Assistant Professor Medicine, Fatima Jinnah Medical University, Lahore, ³Associate Professor Medicine, Fatima Jinnah Medical University, Lahore, ⁴Woman Medical Officer, Mayo Hospital, Lahore, ⁵Radiologist, Sir Ganga Ram Hospital, Lahore, ⁶Assistant Professor, Indiana University, USA/Fatima Jinnah Medical University, Shahrah-e-Fatima Jinnah, Queen's Road, Lahore

Correspondence to: Dr Khadija Muneer, khadijawas@gmail.com

ABSTRACT

Background: Diabetes mellitus is the fifth leading cause of death in developed countries. Risk of infections is increased in diabetic patients with urinary tract being the most common site of infection. Worldwide, the prevalence of urinary tract infections (UTI) is estimated to be around 150 million persons per year. UTI exhibit a poorer prognosis in patients of type 2 diabetes mellitus (T2DM) due to a compromise in their immune status in addition to emerging antimicrobial resistance due to widespread use of antimicrobial agents. This study aims to determine frequency of common bacterial pathogens causing UTI and antibiotic sensitivity pattern among type 2 diabetic patients. This may help to guide physicians regarding appropriate management of UTI in the local population to avoid multidrug resistance.

Patients and methods: This cross sectional study involved 215 type 2 diabetic patients admitted to Medical Unit-III of Sir Ganga Ram hospital, Lahore from 01-12-2016 to 31-05-2017. Type 2 adult diabetic patients (diagnosed for at least 1 year) of age 40-80 years, either male or female, with urine pus cells ≥ 5 /HPF performed as baseline investigation at admission were included in the study. Further urine samples were obtained (clean catch midstream urine 10cc in two wide mouth sterile screw capped plastic jars) before starting treatment. Samples were sent for microscopy and culture. Culture results and antibiotic sensitivities were recorded. Data was analyzed using SPSS for windows version 23.0. Categorical variables like gender, organisms and antibiotic sensitivity was presented in Frequency or Percentage form. Numerical variables like age and duration of diabetes were presented as Mean \pm SD. Data was stratified for age, gender and duration of DM to deal with effect modifiers. Post stratification chi-square test was applied. A-value of ≤ 0.05 was considered significant.

Results: Total of 215 patients were included. Mean age was 63.76 ± 10.342 years of which 164 (76.3%) were females, while 51 (23.7%) were males reflecting female predominance. Most common isolated organism in urine culture was *E. coli* (58.6%), followed by Klebsiella (20.5%), Proteus (9.3%), Pseudomonas (8.4%) and Enterococcus (3.3%) species. Most of the patients (76.7%) were sensitive to piperacillin-tazobactam followed by carbapenems (67%) and Gentamycin (56.7%), whereas 24.2% showed sensitivity to all antibiotics.

Conclusion: Females are most commonly affected among diabetic patients. The commonest isolate was *E. coli* among culture positive UTIs. *E. coli* was most sensitive to piperacillin-tazobactam and carbapenems followed by Gentamycin while least sensitive to Ceftriaxone. The results emphasize the importance of practice of urine culture and sensitivity testing in diabetic patients and cautious approach to use of commonly used antibiotic ceftriaxone and ciprofloxacin as empirical treatment for UTI despite the low sensitivity of the drug to the most prevailing organism.

Keywords: Urinary tract infection; Type 2 Diabetes Mellitus; Antibiotic sensitivity; Pathogens

INTRODUCTION

Diabetes mellitus is one of the most challenging health issues of 21st century considering that it is among the top ten leading causes of mortality in developed countries.¹ Approximately under half a billion people worldwide are suffering from diabetes – global

prevalence of diabetes in 2019 was around 9.3% (463 million people), projected to increase to 10.2% (578 million) by 2030 and 10.9% (700 million) by 2045.² Asian patients not only have a greater risk of developing diabetes but they also exhibit poorer prognosis.³ The prevalence of Diabetes in Pakistan is reported to be around 11.7%.⁴ Type 2 diabetes mellitus (T2DM) is a group of various disorders including several degrees of insulin resistance, improper insulin secretion, and a rise in glucose production. Patients of T2DM have a greater risk of developing infections with the urinary tract infection (UTI) being the most frequent.⁵ The global

Conflict of Interest: The authors declared no conflict of interest exist.

Citation: Fatima K, Muneer K, Hashmat N, Humayun S, Pervaiz Q, Maqsood S. Bacterial urinary tract infection and antibiotic sensitivity pattern among type II diabetic patients – A cross-sectional study from tertiary care teaching hospital in Central Lahore. J Fatima Jinnah Medical Univ. 2019; 13(4): 180-183.

DOI: <https://doi.org/10.37018/jfjmu.684>

prevalence of urinary tract infections acquired in the community is 0.7%.⁶ UTIs are not only more common in type 2 diabetics but the course of disease is also more severe with greater morbidity as a result of a compromised immune system and physiological changes attributable to neuropathies involving the urinary tract.⁴⁻⁷ UTIs in diabetic patients have a range of clinical presentations; from asymptomatic bacteriuria to symptomatic UTI and its complications.⁷ *E. coli* followed by *Klebsiella*, *Pseudomonas*, *Proteus mirabilis* and *Enterococcus* are the most commonly implicated pathogens.⁸ The susceptibility of these organism to antibiotics is variable in different regions of the world.^{9,10} On account of rapidly emerging multidrug resistant (MDR) strains of the organisms to specific antibiotics following noncompliance, injudicious use of drugs, over the counter medications particularly in underdeveloped countries, health care professionals need knowledge of bacterial uropathogens and antibiotic sensitivity patterns in their respective local population regions to avoid emerging resistance to organisms due to irrational use of antibiotics. This study aims to determine frequency of common bacterial pathogens causing UTI and antibiotic sensitivity pattern among type 2 diabetic patients with the objective to guide physicians regarding appropriate management of UTI in the local population to avoid multidrug resistance.

PATIENTS AND METHODS

This cross sectional study was performed in Medical Unit-III of Sir Ganga Ram Hospital, Lahore from 01-12-2016 to 31-05-2017. A sample size of 215 cases was taken by non-probability consecutive purposive sampling using a confidence level of 95% and a margin of error of 4%. UTI is generally asymptomatic in diabetics. Patients of both genders, aged 40- 80 years, with known Type II diabetes diagnosed for at least 1 year and found to have **urine pus cells** ≥ 5 /HPF on baseline urine test at admission were included in the study report. Patients not giving consent, those with urinary catheter, already taking antibiotics or those on immunosuppressive drugs were excluded from the study. A semi-structured close-ended questionnaire was used to collect the demographic and study parameters details. Urine samples were taken using clean catch midstream urine 10cc in two wide mouth sterile screw capped plastic jars before starting treatment. Samples were sent for microscopy and culture to pathology lab of Fatima Jinnah Medical University within an hour of

Table 1. Frequency of UTIs in diabetic patients according to age, gender and duration of diabetes

Characteristics	Frequency	Percentage
Age groups		
40 – 60 years	96	44.7
61 – 80 years	119	55.3
Gender		
Male	51	23.7
Female	164	76.3
Duration of diabetes		
< 15 years	146	67.9
>15 years	69	32.1

sample collection. Samples for culture were inoculated on Blood and Cystine-Lactose-Electrolyte Deficient (CLED) agar media. Cultures showing $>10^5$ CFUs of organisms/ml were taken as positive. For each of the isolates, antibiotic susceptibility was done by Kirby Bauer disk diffusion technique and a record was made of culture results and antibiotic sensitivities. SPSS for windows version 23 was used for data entry and analysis. All the categorical variables like gender, organisms and antibiotic sensitivity were expressed as frequency or percentage. All the numerical variables like age and duration of diabetes were presented as mean \pm SD.

RESULTS

During the study period, 215 patients were included, having mean age of 63.76 ± 10.3 years. Among them 164 (76.3%) were females, while 51 (23.7%) were males. Table 1 summarizes the demographic details of study patients. The mean duration of diabetes recorded in this study sample was 12.71 ± 6.021 years. Out of total 215 patients, 104 (48.4%) patients had *E. coli*, 33 (15.3%) had *Klebsiella*, 19 (8.8%) had *Pseudomonas*, 14 (6.5%) had *Proteus*, 4 (1.9%) had *Enterococcus*, 4 (1.9%) had other species and 37 (17.2%) patients had no growth on urine culture (Table 2). *E. coli* exhibited greatest sensitivity to carbapenems 99% and Piperacillin-tazobactam (99%) followed by, Gentamycin 68.4%, Ciprofloxacin 52.5% and Ceftriaxone 34.5%. *Klebsiella* spp. showed higher sensitivity to carbapenems 99% followed by piperacillin-tazobactam 98%, Gentamycin 75.2%, Ciprofloxacin 68.3% and Ceftriaxone 40.2%.

Table 2. Frequency of microorganism causing UTI

Microorganisms	Number	Percentage
<i>E. coli</i>	104	48.4
<i>Klebsiella</i>	33	15.3
<i>Pseudomonas</i>	19	8.8
<i>Proteus</i>	14	6.5
<i>Enterococcus</i>	4	1.9
No growth	37	17.2
Others (<i>Staphylococcus</i>)	4	1.9
Total	215	100

Table 3. Antibiotic sensitivity pattern of UTI pathogens

Antimicrobial agents	Susceptibility	<i>E. coli</i>	<i>Klebsiella</i>	<i>Pseudomonas</i>	<i>Proteus</i>	<i>Enterococcus</i>
Piperacillin/Tazobactam	Sensitive	99	98	98	98	97
	Resistant	1	2	2	2	3
Carbapenems	Sensitive	99	99	99	98	97
	Resistant	1	1	1	2	3
Gentamycin	Sensitive	68.4	75.2	74.9	73.4	78.1
	Resistant	31.6	24.8	25.1	26.6	21.9
Ciprofloxacin	Sensitive	52.5	68.3	42.8	48.4	2.2
	Resistant	47.5	31.7	57.2	51.6	97.8
Ceftriaxone	Sensitive	34.5	40.2	46.7	68.6	5.5
	Resistant	65.5	59.8	52.3	31.4	94.5

Pseudomonas was highly sensitive to Carbapenems 99% followed by Piperacillin-Tazobactam 98%, Gentamycin 74.9%, Ceftriaxone 46.7% and Ciprofloxacin 42.8%. *Proteus spp.* also showed highest sensitivity to carbapenems 98% and Piperacillin-Tazobactam 98% followed by Gentamycin 73.4%, Ceftriaxone 68.6% and Ciprofloxacin 48.4%. Enterococci had high sensitivity to carbapenems 97% and Piperacillin-Tazobactam 97%, followed by Gentamycin 78.1%, Ceftriaxone 5.5% and Ciprofloxacin 2.2% (Table 2).

DISCUSSION

This study was carried in diabetic patients from Lahore, Pakistan to determine the frequency of UTI and antimicrobial susceptibility pattern. Female diabetic patients showed a greater prevalence of UTI (76%) along with a significant correlation between gender and antibiotic sensitivity. Similar observations were made by Kaleemullah and coauthors in a Pakistani study and Sewify and coworkers from Kuwait.^{8,12} One of the reasons for this increased prevalence reported is the anatomical differences of the urinary tract between males and females (as a shorter urethra allows easier access of bacteria to the bladder) and poor personal hygiene.^{13,14} Majority of patients (55.3%) were in age group of 60-80 years which is in disagreement with other studies which have quoted majority of patients in lesser age groups. In a study from Pakistan¹² majority of patients were in the age group of 50-60 years while in another study from India¹⁴ majority was 56-65 years. This could be due to poor hygiene, lower immunity, social and environmental factors. The commonest isolate was *E. coli* (48.4%) similar to studies from Pakistan, Ethiopia, India and Kuwait while isolates of *Klebsiella* 33 (15.3%), *Pseudomonas* 19 (8.8%), *Proteus* 14 (6.5%), *Enterococcus* 4 (1.9%) and others 4 (1.9%) vary in local and other studies.^{8,12,15-17} The high prevalence of *E. coli* strains can be attributed to their possession of structural virulence factors such as

fimbriae, pili, flagella and secreted virulence factors such as toxins and iron-acquisition systems.¹⁷ *E. coli* exhibited greatest sensitivity to Piperacillin-Tazobactam (99%) and Carbapenems (99%) in this study which is consistent with observation made by Kaleemullah and colleagues where *E. coli* showed a similar sensitivity to Piperacillin-Tazobactam and Imipenem (100%).¹² One study from India showed that organisms were most sensitive to Meropenem.¹⁸ Sensitivity to Piperacillin and Carbapenems was not tested in studies from Ethiopia and Kuwait.^{15,16} Piperacillin/Tazobactam had highest sensitivity towards all uropathogens in this as well as in studies from India and Pakistan.^{12,18,19} In this study most uropathogens showed less sensitivity to Ciprofloxacin and ceftriaxone – the commonly used antibiotics in this region – with *E. coli* showing 52.5% and 34.5% sensitivity respectively. This is consistent with studies from Kaleemullah and associates where *E. coli* was 45.9% sensitive to Ceftriaxone and 38.9% to Ciprofloxacin as well as in studies from India and Ethiopia.^{12,14,15}

The worldwide rise in the incidence of T2DM in the recent years and greater susceptibility of female diabetics to developing UTI with a worse prognosis is expected to place a significant burden on medical costs. This is further complicated with the development of antibiotic resistant strains of UTI pathogens as a result of frequent and improper antibiotic prescription to patients of UTI. There is a clear need for early culture sensitivity in diabetic patients presenting with UTI for prescription of culture specific antibiotics to decrease emergence of resistant organisms, decreasing medical cost and burden on health system resulting from repeated hospitalizations, and avoid complications due to multiple antibiotic regimes.

CONCLUSION

Prevalence of UTI is more among T2DM females. *E. coli* remains the most common organism causing UTIs and most organisms being sensitive to piperacillin-

tazobactam and carbapenems, in contrast to commonly used antibiotics like ceftriaxone and ciprofloxacin, reflecting that antibiotic susceptibilities are changing over time. However further studies are needed to validate these results so as to guide early and judicious use of antibiotics to prevent emerging resistance to routinely used antibiotics.

Acknowledgements: I would like to acknowledge the assistance of Professor Dr. Monazza Gondal, Department of Pathology of Fatima Jinnah Medical University in performing urine analysis and cultures.

REFERENCES

1. IDF Diabetes Atlas 8th edition 2017 [Internet]. Available from: <https://diabetesatlas.org/en/>
2. Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, et al. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Res Clin Pract.* 2019;157:107843.
3. Khan NA, Wang H, Anand S, Jin Y, Campbell NRC, Pilote L, et al. Ethnicity and sex affect diabetes incidence and outcomes. *Diabetes Care.* 2011;34:96-101.
4. Meo SA, Zia I, Bukhari IA, Arain SA. Type 2 diabetes mellitus in Pakistan: Current prevalence and future forecast. *J Pak Med Assoc.* 2016;66(12):1637-42.
5. Nitzan O, Elias M, Chazan B, Saliba W. Urinary tract infections in patients with type 2 diabetes mellitus: review of prevalence, diagnosis, and management. *Diabetes Metab Syndr Obes.* 2015;8:129-36.
6. Tandogdu Z, Wagenlehner FM. Global epidemiology of urinary tract infections. *Curr Opin Infect Dis.* 2016; 29(1):73-9
7. Wan YL, Lee TY, Bullard MJ, Tsai CC. Acute gas-producing bacterial renal infection: correlation between imaging findings and clinical outcome. *Radiology.* 1996;198(2):433-8.
8. Sewify M, Nair S, Warsame S, Murad M, Alhubail A, Behbehani K, et al. Prevalence of Urinary Tract Infection and Antimicrobial Susceptibility among Diabetic Patients with Controlled and Uncontrolled Glycemia in Kuwait. *J Diabetes Res.* 2016;2016:6573215.
9. MergaDuffa Y, TerfaKitila K, MamuyeGebretsadik D, Bitew A. Prevalence and Antimicrobial Susceptibility of Bacterial Uropathogens Isolated from Pediatric Patients at Yekatit 12 Hospital Medical College, Addis Ababa, Ethiopia. *Int J Microbiol.* 2018;2018:8492309.
10. Al- Mazini M. Incidence and sensitivity of bacterial uropathogens among diabetic patients. *Pelagia Research Library European Journal of Experimental Biology,* 2016, 6(3):8-12.
11. Ventola CL. The antibiotic resistance crisis: part 1: causes and threats. *P T.* 2015;40(4):277-83.
12. Zubair KU, Shah AH, Fawwad A, Sabir R, Butt A. Frequency of urinary tract infection and antibiotic sensitivity of uropathogens in patients with diabetes. *Pak J Med Sci.* 2019;35(6):1664-8.
13. Acharya D, Bogati B, Shrestha G, Gyawali P. Diabetes mellitus and Urinary Tract Infection: Spectrum of Uropathogens and their Antibiotic Sensitivity Pattern. *J Manmohan Memorial Inst Health Sci.* 2015;1(4):24-8.
14. Murmu M, K. P. KM, Meher RK, Pradhan B, Kar A, Naik M. Spectrum of uropathogens and their antibiotic sensitivity pattern in diabetes mellitus patients at a tertiary care hospital in Odisha, India. *Int J Res Med Sci.* 2018;6(5):1549-53.
15. Nigussie D, Amsalu A. Prevalence of uropathogen and their antibiotic resistance pattern among diabetic patients. *Turk J Urol.* 2017;43(1):85-92.
16. Woldemariam HK, Geleta DA, Tulu KD, Aber NA, Legese MH, Fenta GM, et al. Common uropathogens and their antibiotic susceptibility pattern among diabetic patients. *BMC Infect Dis.* 2019;19(1):43.
17. Terlizzi ME, Gribaudo G, Maffei ME. UroPathogenic Escherichia coli (UPEC) Infections: Virulence Factors, Bladder Responses, Antibiotic, and Non-antibiotic Antimicrobial Strategies. *Front Microbiol.* 2017;8:1566.
18. Kiranmala K, Johnson R, Savio J, Idiculla J. Microbiologic profile and clinical practices in urinary tract infections in a tertiary care center in Southern India. *J Family Med Prim Care* 2019;8:2888-92.
19. Bashir H, Saeed K, Jawad M. Causative agents of urinary tract infection in diabetic patients and their pattern of antibiotic susceptibility. *KMUJ.* 2017;9(4):201-4.